

PEST CONTROL OUTDOORS: FIRE ANTS

General Information

GENERAL INFORMATION

D-FENSE SC is a suspension concentrate insecticide for use both indoors and outdoors for effective knockdown and long residual control of listed pests.

For Use in Residential, Industrial and Institutional Areas

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

RESTRICTIONS:

1. DO NOT apply D-FENSE SC through any type of irrigation system.
2. DO NOT water the treated area to the point of run-off.
3. DO NOT make applications during rain.
4. DO NOT apply this product to edible crops.
5. DO NOT apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm.
6. DO NOT apply in classrooms when in use.
7. DO NOT apply this product in the immediate area when occupants are present (including people and animals).
8. Do not apply this product more than once every 21 days. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - a. Treatment to soil or vegetation around structures;
 - b. Applications to lawns, turf, and other vegetation;

c. Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow this product to enter any drain during or after application.

Limitations, Restrictions, and Exceptions

PEST CONTROL OUTDOORS: FIRE ANTS

Dilute 1.5 oz. per one gallon of water and apply D-FENSE SC as a drench treatment. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding area to a 2-foot diameter. Best results are obtained when applied in cool weather, 65 to 80°F or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.

Method

[Drench](#)

Rates

[field rates 0](#)

•

Timings

[Treat new mounds as they appear](#)

[In cool weather, 65 to 80°F or in early morning or late evening hours](#)