

# **APPLES, PEARS, PEACHES, CHERRIES, PLUMS, GRAPES**

## General Information

### APPLICATION INSTRUCTIONS

NOTE: All rates are calculated for mature trees with normal foliage cover. Rates should be adjusted downward proportionally for smaller immature trees. All applications should be made in sufficient water to insure thorough coverage without significant runoff. Minimum GPA for concentrate sprays is 10 to 20 for aerial application and 10 gallons for concentrate ground application. Dilute sprays may be applied at a minimum of 60 GPA, and a maximum of 1000 gallons. When tank mixing 435 oil with other pesticides, check compatibility (jar-test) before mixing in application equipment. Follow all precautions and instructions on all pesticide labels being tank mixed.

### Limitations, Restrictions, and Exceptions

APPLES, PEARS, PEACHES, CHERRIES, PLUMS, GRAPES: Use 1/2 cup to one gallon of (1.5 gal./50 gal. of water) to control European Red mite eggs, certain aphids, and scale insects. Spray thoroughly, wetting insects on both sides of leaves.

PRECAUTIONS: Do not spray within 30 days before or after using Sulfur or Captan. Do not apply when temperatures are above 95 °F. Do not apply oil when rain or snow is likely to follow quickly or when the temperature is below 40 degrees F., or is likely to drop to the freezing point before the spray has thoroughly dried. Use the proper amounts only as stated on the label and follow all cautions.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

### Restricted Entry Interval

4 hours

Timings

[N.A.](#)