

# ORNAMENTAL PLANTS - AZALEAS AND CAMELLIAS

## General Information

### APPLICATION INSTRUCTIONS

NOTE: All rates are calculated for mature trees with normal foliage cover. Rates should be adjusted downward proportionally for smaller immature trees. All applications should be made in sufficient water to insure thorough coverage without significant runoff. Minimum GPA for concentrate sprays is 10 to 20 for aerial application and 10 gallons for concentrate ground application. Dilute sprays may be applied at a minimum of 60 GPA, and a maximum of 1000 gallons. When tank mixing 435 oil with other pesticides, check compatibility (jar-test) before mixing in application equipment. Follow all precautions and instructions on all pesticide labels being tank mixed.

### Limitations, Restrictions, and Exceptions

PRECAUTIONS: Do not use more than 1 gallon per 100 gallons of spray, as injury may result under certain circumstances. Do not use oil sprays on foliage within 30 days before or after using Sulfur or Captan. Do not spray broadleaved evergreens when temperatures are near or below freezing. Dormant oil spray may cause discoloration of Blue Spruce. Do not repeat oil sprays in less than 30 days of first applications, nor use more than twice in twelve month period. Do not spray trees or plants weakened from any cause whatsoever such as lack of moisture, excess moisture, cold or freeze damage, recent root pruning or transplanting. Avoid spraying trees or plants when temperatures are excessively high or low. Keep guns or spray wand in motion while spraying. Do not overlap with additional spray onto foliage already dry.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

### Restricted Entry Interval

4 hours

### Timings

[Apply in spring after blooming and in fall prior to blooming, and at least 3 weeks](#)

before anticipated freeze.