

ESTABLISHED ORNAMENTALS IN RESIDENTIAL AREAS - FOLIAR APPLICATIONS

Limitations, Restrictions, and Exceptions

ESTABLISHED ORNAMENTALS IN RESIDENTIAL AREAS

Foliar Applications

Use Dominion Tree & Shrub as a foliar spray to provide contact control of larvae, nymphs or adults of the following insect pests: adelgids, aphids, Japanese beetles, lace bugs, leaf beetles (including elm leaf beetles and viburnum leaf beetles), leafhoppers (including glassywinged sharpshooter), mealybugs, psyllids, sawfly larvae, thrips (suppression) and whiteflies.

Application Instructions

Apply when insects first appear and before high pest populations are established. Retreat when re-infestation occurs, when high pest pressure is observed or repeat applications on an as needed basis with 10-14 day intervals between applications.

How to Apply

Foliar applications with water must be made as a thorough cover spray to provide uniform distribution. Spray to wet all foliage (leaves, stems and branches). Target the undersides of the leaves and try to penetrate dense foliage with the spray droplets.

Hand-Operated Sprayers

Including backpack, compression, knapsack or tank type sprayers.

For foliar applications mix 1.5 (1 1/2) teaspoons (tsp) of Dominion Tree & Shrub per one (1) gallon of water.

Hose-End Sprayer Instructions [For Gilmour Dial-A-Mix Sprayer or Air-O-Matic]

1. Set the dial to 1 1/2 teaspoons (tsp.). Keep the dial set to this setting while spraying.
2. Add an adequate amount of product to the spray bottle to treat the total area. Do not add water to jar.
3. Spray to wet all foliage and direct the spray droplets for an even distribution on the target plants. Any unused product can be poured back into its original container.

Method

[Foliar spray](#)

Rates

[field_rates 0](#)

•

Timings

[When insects first appear and before high pest populations are established.](#)