

GRASS AND DICHONDRA LAWNS - GRUBS, JAPANESE BEETLE GRUBS, ATAENIUS SPRETULUS (DUNG BEETLES)

General Information

DIRECTIONS FOR USE

Three Easy Steps to Use this Product

Step #1: Determine size of the job, measure the area to be treated.

Step #2: Use spreader setting chart to determine appropriate setting for your spreader.

Step #3: Treat as directed in rate chart. Thorough coverage is important. Water in granules where indicated.

Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

Limitations, Restrictions, and Exceptions

GRASS AND DICHONDRA LAWNS

Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4 - 1/2 inches of water after application. Treat again in late summer or early fall if a second generation occurs.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

-

Timings

[Billbug: When activity is first observed or when chewed or brown grass indicates](#)

damage from these insects.

Grub (except dung beetle): Treat any time between late July and early October.

Dung beetle: In spring or early summer when damage is first noticed.