

APPLICATIONS IN FOOD PROCESSING ESTABLISHMENTS

General Information

PRODUCT INFORMATION

PYROFOS CS is a liquid microencapsulated concentrate containing the active ingredient Chlorpyrifos for use in and around (within 6 to 10 feet) the following sites:

Calf Hutches and Calving Pens and Parlors; Cargo Holds of Ships, Empty Chicken Houses, Food and Non-Food Areas of Manufacturing, Industrial Plants and Food Processing Plants; Hog Barns; Milk rooms; Railroad Boxcars.

PYROFOS CS may be used to treat any surface which will not be damaged or marked by water. Heavy applications may leave a noticeable deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

PYROFOS CS Controls: Angoumois Grain Moths, Aphids, Ants(except carpenter, fire, harvester, and pharaoh), Bees, Beetles, Boxelder Bugs, Cadelles, Carpet Beetles, Cigarette Beetles, Clover Mites, Cluster Flies, Confused Flour Beetle, Crickets, Darkling Beetles, Dermestid Beetles, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Elm Leaf Beetles, Firebrats, Flea Beetles, Flies, Gnats, Grain Mites, Indian meal Moths, Leafhoppers, Leaf Miner, Mealworms, Mediterranean Flour Moths, Millipedes, Moths, Northern Fowl Mites, Psocids, Rod Flour Beetle, Saw-toothed Grain Beetles, Scale, Silverfish, Sowbugs, Spiders(except black widow and brown recluse), Stink Bugs, Trogoderma, and Weevils (Granary Weevils and Rice Weevils).

Limitations, Restrictions, and Exceptions

APPLICATIONS IN FOOD PROCESSING ESTABLISHMENTS

FOR USE IN: Food and Non-Food Areas of Manufacturing, Industrial and Food Processing Plants. Non-food areas include boiler rooms, garbage rooms, drains (to sewers), entries and vestibules, machine rooms, garages, mop closets, and storage (after canning or bottling). Treating within food areas of food processing plants is limited to spot and/or crack and crevice treatments.

Treat where insects are found or normally occur. Application of a 0.25% solution is

suitable for control of light pest infestations (i.e. maintenance

applications). Applications of a 0.5% solution provides residual control of heavy infestations (i.e. clean-out applications) and where extended residual control is desired. A period of 4-7 days is usually required to observe maximum effectiveness on insect populations. Do not re-apply more than once per every 7 days. Apply spray solution so that spray drift is minimized to avoid non-target surfaces. When using as a spot treatment, do not exceed 2 sq. ft. per individual spot.

Method

[Spot treatment](#)

Rates

[field rates 0](#)

-

Timings

[N. A.](#)