

PLANTAIN LILY - FOR PROMOTION OF LATERAL GROWTH ON FINISHED PLANTS

General Information

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

FOR USE IN: COMMERCIAL GREENHOUSES, GLASSHOUSES (WHERE PLANTS ARE GROWN IN CONTAINERS), AND ON ACTIVELY GROWING STOCK PLANTS.

CONFIGURE increases lateral or basal branching and promotes flowering in certain ornamental crops. In addition, applications of CONFIGURE reduce the overall height of the plant resulting in more compact and marketable plants.

Plant response to CONFIGURE is strongly influenced by cultural and environmental variables. Growing media, water/fertilizer management, temperature, light, greenhouse composition, and other cultural practices may impact plant response. Plant size, container size and cultivar or variety can also influence treatment response. Although multiple cultivars have been tested for sensitivity to CONFIGURE and have demonstrated high levels of tolerance, it is impossible to insure an acceptable response in all cultivars. First time users of CONFIGURE should first conduct trials on a limited number of plants to determine crop response.

APPLICATION INFORMATION

Apply CONFIGURE as a foliar spray using standard foliar application spray equipment. Make sure that sufficient volumes are used to thoroughly wet foliage. Spray uniformity is equally important. Uniformly apply 1-2 quarts of finished spray solution to 100 sq. ft. of area. A high quality wetting agent or spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions.

Refer in the label for CONFIGURE DILUTION GUIDE.

Limitations, Restrictions, and Exceptions

DIRECTIONS FOR USE

Plantain Lily (Hosta spp.):

Use CONFIGURE on Hosta to promote lateral growth of finished plants by inducing the outgrowth of axillary and rhizomic buds. Use CONFIGURE to increase offsets during propagation.

For the promotion of lateral growth on finished plants: Apply CONFIGURE at rates ranging from 1000 - 3000 ppm in a uniform spray volume. CONFIGURE is most effective when plants are fully established prior to application (i.e., at least 3 to 4 weeks after potting), when there is evidence of surface root development, but before flower initiation.

Method

[Foliar spray](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[When there is evidence of surface root development, but before flower initiation.](#)