

# **TABLE GRAPES (SEEDLESS) - RAISIN (SIZING) - 3 TO 5 TARGET DIAMETER**

## General Information

### GENERAL INFORMATION

- Use only as directed. Read thoroughly and understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See Conversion Table at end of Directions for Use section.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of this label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Ensure that the pH of the water is less than 8.5.
- Absorption of FALGRO 4L into the plant is greatest under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying. Re-apply FALGRO 4L if significant rain occurs within 2 hours of application.
- FALGRO 4L has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

## Limitations, Restrictions, and Exceptions

### SPRAYING GUIDE FOR GRAPES

Application to grapes is best made by ground sprayer. Spray volume, quality and direction are chosen to ensure thorough wetting of all flower clusters and berries.

## TABLE GRAPES

Application of FALGRO 4L for 'Sizing' produces larger berries and clusters when used in conjunction with girdling and thinning techniques.

Timing: 1-4 sprays starting at these berry sizes. All applications to be made within 14 days.

Method

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[1-4 sprays starting at these berry sizes.](#)