

GRAPEFRUIT (ALL APPROVED STATES EXCEPT CALIFORNIA) - TO DELAY RIND AGEING AND PREVENT PRE-HARVEST DROP OF MATURE FRUIT AND INCREASES PEEL STRENGTH

General Information

GENERAL INFORMATION

- Use only as directed. Read thoroughly and understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See Conversion Table at end of Directions for Use section.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of this label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Ensure that the pH of the water is less than 8.5.
- Absorption of FALGRO 4L into the plant is greatest under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying. Re-apply FALGRO 4L if significant rain occurs within 2 hours of application.
- FALGRO 4L has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

SPRAYING GUIDE FOR CITRUS

Spray volume, quality and direction are chosen to ensure thorough wetting of all fruit. Do not exceed maximum application rates. Do not apply to trees of low vigor

or suffering stress of any kind as this can cause severe fruit and/or leaf drop. Some drop of older leaves may occur after application of gibberellic acid. Never tank mix FALGRO 4L with materials creating a high pH in the tank, e.g., white wash. The use of copper fungicides and/or oils within three weeks of a FALGRO 4L application may result in significant leaf and/or fruit drop.

GRAPEFRUIT (All approved states except California)

All Grapefruit Varieties Except Star Ruby:

To delay rind ageing and prevent pre-harvest drop of mature fruit and increases peel strength: Spray 16 to 56 g ai/acre as a dilute spray in a minimum of 250 gallons of water per acre. Ensure thorough wetting. Application of FALGRO 4L also reduces water loss during storage and can be used to manage harvest timing.

Early spray (Before color change)

Make one application 14 days before color break (usually August to September). Application at this timing produces the greatest delay in rind ageing and the firmest rind possible.

Late spray (After color break)

Make one application after marketable color has developed (usually October to December). Late application may cause some re-greening of the fruit.

Do not apply to areas that are destined to be harvested early as treatment will delay color change. Application after coloring will cause fruit to re-green if left on the tree for an extended period of time. Avoid applications after December or after trees break dormancy in order to avoid adversely affecting the next crop. Be aware that the efficacy of treatment varies from season to season, depending on environmental conditions.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

Early spray: Before color change.

Late spray: After color break.