

VALENCIA ORANGES AND OTHER ROUND ORANGES - TO DELAYED RIND AGEING AND SOFTENING

General Information

GENERAL INFORMATION

- Use only as directed. Read thoroughly and understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See Conversion Table at end of Directions for Use section.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of this label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Ensure that the pH of the water is less than 8.5.
- Absorption of FALGRO 4L into the plant is greatest under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying. Re-apply FALGRO 4L if significant rain occurs within 2 hours of application.
- FALGRO 4L has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

SPRAYING GUIDE FOR CITRUS

Spray volume, quality and direction are chosen to ensure thorough wetting of all fruit. Do not exceed maximum application rates. Do not apply to trees of low vigor or suffering stress of any kind as this can cause severe fruit and/or leaf drop. Some

drop of older leaves may occur after application of gibberellic acid. Never tank mix FALGRO 4L with materials creating a high pH in the tank, e.g., white wash. The use of copper fungicides and/or oils within three weeks of a FALGRO 4L application may result in significant leaf and/or fruit drop.

VALENCIA ORANGES (All approved states) & OTHER ROUND ORANGES (All approved states except California)

To delay rind ageing and softening: Make one application of 40 to 80 g ai/acre as a dilute spray or concentrate in August to October.

This application is expected to delay color development and re-green any mature fruit present at application. After marketable color has been achieved, beneficial treatment effects will be reduced the longer the fruit remains on the tree.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[August to October.](#)