

STATICE (ALL APPROVED STATES EXCEPT CALIFORNIA)

General Information

GENERAL INFORMATION

- Use only as directed. Read thoroughly and understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See Conversion Table at end of Directions for Use section.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of this label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Ensure that the pH of the water is less than 8.5.
- Absorption of FALGRO 4L into the plant is greatest under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying. Re-apply FALGRO 4L if significant rain occurs within 2 hours of application.
- FALGRO 4L has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

STATICE (All approved states except California)

Application of FALGRO 4L promotes early flowering and increases flower yield. Make one application of 40 – 50 g ai in 25 gallons of water when plants are more than 10 inches in diameter (approx. 90 – 110 days after planting). Apply 10 ml of spray as a drench to each plant. Do not exceed the specified dose or repeat treatment. Initiation of flowering is influenced by extended photoperiod, nutrition and reduced

night temperature. Treatment with FALGRO 4L lessens the need for reduced night temperature and long photo-period.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[When plants are more than 10 inches in diameter \(approx. 90-110 days after planting\).](#)