

CITRUS AND AVOCADOS

General Information

GENERAL INFORMATION

Hye-Green 2-18-40 is a water-soluble foliar fertilizer containing various micronutrients important to healthy plant development. It should be mixed with water prior to application and may be used in conjunction with pesticide sprays except lime sulfur and highly alkaline spray materials.

The rates given under DIRECTIONS FOR USE should be adjusted according to the crop stage of development, the number of applications intended, and soil and weather conditions. If only a single application is intended or when soil and weather conditions are unfavorable for crop growth, higher rates are recommended. Lower rates are recommended when leaf area is small, new growth exists or on new plantings. For best results, applications should begin as soon as the leaves are large enough to absorb and utilize the nutrients and should continue at 10-14 day intervals during the primary growth period.

Hye-Green 2-18-40 can be used in crop protection sprays with a minimum volume of 20 gallons of spray per acre by ground and 5 gallons of spray per acre by air; however, the concentration of Hye-Green 2-18-40 solution should not exceed 20 pounds in 100 gallons of water. Avoid spraying during full bloom; except citrus. When spraying blossoms, use lower rates. Do not store under excessive hot or cold temperatures for long periods. May be corrosive to mild steel and brass.

Limitations, Restrictions, and Exceptions

CITRUS AND AVOCADOS

- Apply Hye-Green 2-18-40 at 5-20 pounds per acre in sufficient water to cover.
- Make applications at pre-bloom stage, when fruit is sizing, and after harvest.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

•

Timings

[Post-harvest](#)

[Pre-bloom stage.](#)

[When fruit is sizing.](#)