

# **CITRUS, FIELD, ROW, VEGETABLE, FRUIT, TREE, VINE, ORNAMENTAL AND TURF CROPS - FOLIAR APPLICATION (DILUTE)**

## General Information

### GENERAL INFORMATION

LIQUID ZINC 9% is especially recommended for citrus crops but may also be used on field, row, vegetable, fruit, tree, vine, ornamental and turf crops. LIQUID ZINC 9% alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. LIQUID ZINC 9% may be applied via ground sprayer (high-volume or low-volume), aircraft or drip irrigation systems.

### DIRECTIONS FOR USE

Soil applications of LIQUID ZINC 9% are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliar applied LIQUID ZINC 9% most efficiently. Immediate post-irrigation or post-rainfall application gives the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of LIQUID ZINC 9%.

### Limitations, Restrictions, and Exceptions

#### CITRUS, FIELD, ROW, VEGETABLE, FRUIT, TREE, VINE, ORNAMENTAL AND TURF CROPS

LIQUID ZINC 9% applications should be a minimum of 7-14 days apart. Apply 2-4 quarts/acre as needed on a 7-14 day schedule. AVOID foliar applications of LIQUID ZINC 9% when plants are under moisture stress. AVOID foliar applications of LIQUID ZINC 9% to deciduous tree crops during bloom.

Dilute Applications: Apply the suggested dose of product in enough water for thorough coverage.

### Method

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

- 

Timings

[N.A.](#)