

MOLES

General Information

DIRECTIONS FOR USE

USE RESTRICTIONS: This product may be used only to control eastern moles (*Scalopus aquaticus*), star-nose moles (*Condylura cristata*), hairy-tailed moles (*Parascalops breweri*), coast mole (*Scapanus orarius*), broad-footed mole (*S. latimanus*) or Townsend's moles (*S. townsendii*) on lawns, turf areas, golf courses, and other non-food grassy areas. Bait must be applied directly into main underground tunnel or subsurface runways. Do not place bait above the ground surface.

SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of surface ridges in the turf or by a series of conical mounds of earth pushed up from deep burrows. Conical mounds may betray the location of main underground runway.

BAITING: Prior to treatment, determine which burrow systems are active. Using a round-ended wood rod, such as a broomstick, probe ground in vicinity of conical mounds or surface ridges until main underground runway is detected by a sudden decrease in resistance against the probe. Mark open burrow systems and revisit them in 2-3 days. For deeper tunnels, open system with a shovel and leave open. Burrow systems that are active are likely to be plugged by moles.

Treat only active burrow systems. To treat a burrow system, locate main runways by probing. Remove the cap from the tip of the applicator syringe and turn dosing ring to mark indicating half ounce. Inject half of an ounce of bait (one-sixth of the contents of one syringe) through probe or shoveled opened hole at each bait placement location. Make 6 such placements per burrow system. Cover holes made to enter tunnel with sod and soil, but do not allow dirt to sift in to foul or cover the bait.

Check effects of treatment 4-5 days after bait application by opening and marking burrows in the manner used to check for initial activity. Return 2-3 days later and retreat all active burrows. Maximum application rate is 1 syringe/10,000 sq ft.

Limitations, Restrictions, and Exceptions

SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of surface ridges in the turf or by a series of conical mounds of earth pushed up from deep burrows. Conical mounds may betray the location of main underground runway.

BAITING: Prior to treatment, determine which burrow systems are active. Using a round-ended wood rod, such as a broomstick, probe ground in vicinity of conical mounds or surface ridges until main runway is detected by a sudden decrease in resistance against the probe. Mark open burrow systems and revisit them in 2-3 days. For deeper tunnels, open system with a shovel and leave open. Burrow systems that are active are likely to be plugged by moles.

Treat only active burrow systems. To treat a burrow system, locate main runways by probing. Remove the cap from the tip of the applicator syringe and turn dosing ring to mark indicating half ounce. Inject half of an ounce of bait (one-sixth of the contents of one syringe) through probe or shovel opened hole at each bait placement location. Make six (6) such placements per burrow system. Cover holes made to enter tunnel with sod and soil, but do not allow dirt to sift to foul or cover the bait.

Check effects of treatment 4-5 days after bait application by opening and marking burrows in the manner used to check for initial activity. Return 2-3 days later and retreat all active burrows.

Maximum application rate is 1 syringe/10,000 ft².

Method

[Directed](#)

[Bait](#)

Timings

[N.A.](#)