

DIRECTIONS FOR INCUBATORS AND HATCHERS USING FOGGING DEVICES

General Information

Spectrasol Disinfectant is a phosphate-free, pH neutral formulation designed to provide effective cleaning, deodorizing, and disinfection for hospitals, nursing homes, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, athletic facilities, sports stadiums, amphitheaters and convention centers where housekeeping is of prime importance in controlling cross-contamination from treated surfaces.

Spectrasol Disinfectant, when used as directed, is formulated to disinfect the following hard non-porous inanimate environmental surfaces: Floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, artificial turf surfaces, vanity tops, shower stalls, bathtubs, and cabinets.

Spectrasol Disinfectant is for use in:

- Food processing plants, dairy farms, hog farms, poultry and turkey farms and egg processing plants.
- Veterinary clinics, animal life science laboratories, kennels and cages, breeding and grooming establishments, pet animal quarters, zoos, pet shops, and other animal care facilities.
- For larger areas such as operating rooms and patient care facilities, this product is designed to provide both general cleaning and disinfecting.

Spectrasol Disinfectant is a one-step neutral disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, HIV-2, HBV and HCV) and inhibits the growth of mold and mildew and their odors when used as directed.

Limitations, Restrictions, and Exceptions

Directions for Incubators and Hatchers Using Fogging Devices: Mix 12 ounces of Spectrasol Disinfectant to 122 ounces of water. Fog 3-8 ounces of this into setters

and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breathe this fog and do not enter until the fog has settled (30-60 minutes after fogging is completed). It is acceptable to fog setters and hatchers with a one (1) ounce per every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. Thoroughly scrub all treated feed racks, mangers, troughs, automatics feeders, fountains and waterers with soap and detergent, and rinse with potable water before reuse.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

Method

[Fogging](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[Apply immediately after transfer.](#)