

FIGS

General Information

GENERAL INFORMATION

Apply this product only as specified on the label. Do not use this product for any uses other than those specified on the label. Yellow Jacket 80 DF is formulated for use in water as a spray on fruit, vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined in the spray tank with most commonly used fungicides and insecticides. Pour the specific amount of this product on surface of water in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear type spray equipment. Do not combine with emulsified liquids unless previous use of the mixture has proven physically compatible and safe to plants.

NOTE: Do not use sulfur in combination with oil or within 21 days of an oil spray. Do not use sulfur on apples or other sensitive crops when shade temperatures exceed, or are likely to exceed 90° F. Do not apply or allow sulfur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tung. Do not apply to or allow spray drift on sulfur-sensitive plants such as D'anjou pears, apricots, chrysanthemums, cranberries, spinach, or other sensitive crops listed on the labeling. To do so may result in severe fruit and leaf injury. When growing crops for processing, consult the processor before applying sulfur.

DILUTE GROUND APPLICATION A - Orchard application -Apply specified rate in 100-800 gallons of water per acre. B - All other applications -Apply specified rate in 20-60 gallons of water per acre.

CONCENTRATE GROUND APPLICATION A - Orchard application -Apply specified rate in 20-100 gallons of water per acre. B - All other applications -Apply specified rate in not less than 5 gallons of water per acre.

AERIAL APPLICATION A - Orchard application -Should not be applied in less than 10 gallons of water per acre. B - All other applications -Apply specified rate in 5-20

gallons of water per acre.

Recommendations: Pour needed amount into spray tank under agitation. If necessary, add a few drops of oil to avoid foaming.

Limitations, Restrictions, and Exceptions

FIGS

Since the effectiveness of sulfur varies in different localities, State Agricultural Experiment Stations should be consulted as to the effectiveness before application.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[When mites first appear.](#)