

VOLE CONTROL IN NURSERIES, TREE AND FORESTRY PLANTATIONS, CHRISTMAS TREE FARMS, AND BORDER AREAS / BUFFER STRIPS ADJACENT TO CROPS - HAND SPOT BAITING

Limitations, Restrictions, and Exceptions

VOLE CONTROL IN NURSERIES, TREE AND FORESTRY PLANTATIONS, CHRISTMAS TREE FARMS, AND BORDER AREAS/BUFFER STRIPS ADJACENT TO CROPS:

USE RESTRICTIONS:

For control of only voles in commercial nurseries, tree and forestry plantations, Christmas tree farms, and border areas and buffer strips adjacent to crops (within 100 feet of the edge of the cropland). This product must not be applied directly to food or feed crops except as specified above. Do not apply within 50 feet of any body of surface water or where raptors are actively feeding on voles. Do not allow animals to graze in treated areas. Do not use hay cut after application for feed or bedding. To avoid exposing nontarget organisms, follow the instructions in the \"Pesticide disposal\" section to insure proper clean up of any bait for reuse or disposal. Apply only by hand spot baiting and ground broadcasting. Do not apply aerially. In Colorado, Florida, New York and Vermont, only apply by \"Hand Spot Baiting.\" In Florida, only use this product where Pine voles are known to occur and only apply by \"Hand Spot Baiting.\"

APPLICATION DIRECTIONS:

Before application, locate vole trails, runway systems and harborage areas to be treated. Hand spot baiting: Place 1-1/2 ounces (6 tablespoons) of bait in each active hole, trail or runway (do not exceed 10 lbs per acre). Cover each placement with grass or shingle to avoid exposing nontarget organisms, or place bait in a tamper resistant bait station.

Method

[Spot treatment](#)

[Bait](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[N.A.](#)