

FOLIAR APPLICATION FOR ALL CROPS - DILUTE APPLICATIONS

General Information

GENERAL INFORMATION

LIQUID MAGNESIUM 4% is recommended to prevent and correct magnesium deficiencies of field, row, vegetable, fruit, tree, vine, ornamental and turf crops. LIQUID MAGNESIUM 4% alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. LIQUID MAGNESIUM 4% may be applied via ground sprayers (high-volume or low-volume), aircraft or fertigation systems.

DIRECTIONS FOR USE

Soil applications of LIQUID MAGNESIUM 4% are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly applied LIQUID MAGNESIUM 4% most efficiently. Immediate post-irrigation or post-rainfall application give the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of LIQUID MAGNESIUM 4%.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATIONS FOR ALL CROPS:

Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

Dilute Applications: For rates higher than 2 quarts per acre, split applications of 2 quarts or less are preferred to minimize phytotoxicity.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[N. A.](#)