

FOLIAR APPLICATION FOR ALL CROPS - AERIAL APPLICATIONS

General Information

GENERAL INFORMATION

LIQUID MANGANESE 7% is recommended to prevent and correct manganese deficiencies of field, row, vegetable, fruit, tree, vine, ornamental and turf crops. LIQUID MANGANESE 7% alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. LIQUID MANGANESE 7% may be applied via ground sprayers (high-volume or low-volume), aircraft or fertigation systems.

DIRECTIONS FOR USE

Soil applications of LIQUID MANGANESE 7% are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly applied LIQUID MANGANESE 7% most efficiently. Immediate post-irrigation or post-rainfall application give the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of LIQUID MANGANESE 7%.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATION FOR ALL CROPS

Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

Aerial Applications: Up to 1 quart of LIQUID MANGANESE 7% per 5 gallons of spray may be used, but more dilute solutions will improve coverage and results.

Method

[Broadcast/Foliar Air](#)

Rates

field_rates 0

-

Timings

N. A.