

SOIL APPLICATION FOR ALL CROPS - MILD DEFICIENCY

General Information

GENERAL INFORMATION

LIQUID PISTACHIO MIX is especially recommended to prevent and correct copper, magnesium and zinc deficiencies in pistachios, but it may also be used on field, row, vegetable, fruit, nut, and vine crops. LIQUID PISTACHIO MIX alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. LIQUID PISTACHIO MIX may be applied via ground sprayers (high-volume or low-volume), aircraft or fertigation systems.

DIRECTIONS FOR USE

Soil applications of LIQUID PISTACHIO MIX are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly applied LIQUID PISTACHIO MIX most efficiently.

Immediate post-irrigation or post-rainfall application give the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of LIQUID PISTACHIO MIX.

Limitations, Restrictions, and Exceptions

SOIL APPLICATION FOR ALL CROPS

The lower rates are for field, row and vegetable crops. The higher rates are for fruit, tree, vine, ornamental and turf crops.

Fertigation: Apply the recommended amount of product in split applications one to two weeks apart throughout the growing season.

Method

[Soil application](#)

Rates

[field rates 0](#)

-

Timings

[N. A.](#)