

SOIL APPLICATION FOR ALL CROPS - SEVERE DEFICIENCY

General Information

General Information

BRANDT ORGANICS 10% Zn is recommended to prevent and correct zinc deficiencies of field, row, vegetable, fruit, tree, and vine crops. Applied alone, with pesticides, or with other fertilizers, this product may be applied directly to the soil or to the foliage of growing crops via ground sprayers, aircraft or fertigation systems.

Mixing and Handling Instructions

Agitate before use. Fill spray tank with half of required water volume and begin agitation. Add products in this order, mixing thoroughly after each addition: adjuvants, pesticides, then fertilizers. Fill tank with remainder of water and continue agitation until spray solution is completely mixed. Check chemical mixture compatibility using a jar test prior to application.

Limitations, Restrictions, and Exceptions

Soil Application: The lower rates are for field, row and vegetable crops. The higher rates are for fruit, tree, and vine crops.

Fertigation: Apply the recommended amount of product in split applications 1-2 weeks apart throughout the growing season.

Optimum rate of application will vary depending on treatment interval, soil properties (such as pH, organic matter content, texture), weather conditions, time of year, plant species and its nutrient requirements. For best results, follow soil/tissue test recommendation.

Method

[Soil application](#)

Rates

[field rates 0](#)

-

Timings

N.A.