

## **FOLIAR APPLICATION - OTHER CROPS**

### General Information

#### General Information

BRANDT ORGANICS 10% Zn is recommended to prevent and correct zinc deficiencies of field, row, vegetable, fruit, tree, and vine crops. Applied alone, with pesticides, or with other fertilizers, this product may be applied directly to the soil or to the foliage of growing crops via ground sprayers, aircraft or fertigation systems.

#### Mixing and Handling Instructions

Agitate before use. Fill spray tank with half of required water volume and begin agitation. Add products in this order, mixing thoroughly after each addition: adjuvants, pesticides, then fertilizers. Fill tank with remainder of water and continue agitation until spray solution is completely mixed. Check chemical mixture compatibility using a jar test prior to application.

#### Limitations, Restrictions, and Exceptions

Foliar Application: Actively growing plants with young leaves absorb foliarly applied BRANDT ORGANICS 10% Zn most efficiently. Immediate post-irrigation or post-rainfall application will enhance foliar uptake. Thorough coverage and wetting are needed for optimum foliar spray results.

Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

Dilute: Other Crops: For rates higher than 2 quarts per acre, split applications of 2 quarts or less are preferred to minimize phytotoxicity.

#### Method

[Broadcast/Foliar Air](#)

[Foliar spray](#)

#### Rates

[field\\_rates 0](#)

- 

Timings

[N.A.](#)