

MILD DEFICIENCY

General Information

GENERAL INFORMATION:

LIQUID ZINC 10% is recommended to prevent and correct zinc deficiencies of field, row, vegetable, fruit, tree, vine, ornamental and turf crops. LIQUID ZINC 10% alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. LIQUID ZINC 10% may be applied via ground sprayers (high-volume or low-volume), aircraft or fertigation systems.

DIRECTIONS FOR USE:

Soil applications of LIQUID ZINC 10% are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly applied LIQUID ZINC 10% most efficiently. Immediate postirrigation or post-rainfall application give the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of LIQUID ZINC 10%.

Limitations, Restrictions, and Exceptions

SOIL APPLICATIONS FOR ALL CROPS*

Mild Deficiency 2-4 qts./acre

*The lower rates are for field, row and vegetable crops. The higher rates are for fruit, tree, vine, ornamental and turf crops.

Method

[Soil application](#)

Rates

[field rates 0](#)

-

Timings

N.A.