

## **FOLIAR APPLICATION: TREES (CITRUS)**

### General Information

#### GENERAL INFORMATION

Brandt Sequestar 14% Cu is a fully chelated, water soluble, powdered form of copper EDTA and is recommended for the prevention and correction of copper deficiency in crops. The preferred use is as a soil application to prevent the deficiency although it may also be used as a foliar spray to provide correction when a soil application is impractical.

#### Limitations, Restrictions, and Exceptions

Foliar Application: BRANDT SEQUESTAR 14% Cu may be applied in water or in combination with most pesticides. After the correct dilution has been made, the spray solution should be buffered to pH 6-6.5. Thorough coverage and the use of wetting agents often enhance nutrient uptake from foliar sprays. Application rates and dilution factors depend on crop sensitivity, the amount of foliage to be sprayed and the application method. Avoid applying this product when plants are suffering from moisture stress. If there is any doubt, apply the spray solution to a small test area of the crop or foliage to assess any undesirable effects or phytotoxicity before general application.

Apply any time except during bloom.

Brandt Sequestar 14% Cu may be applied in water or in combination with most pesticides. After the correct dilution has been made, the spray solution should be buffered to pH 6-6.5. Thorough coverage and the use of wetting agents often enhance nutrient uptake from foliar sprays. Application rates and dilution factors depend on crop sensitivity, the amount of foliage to be sprayed and the application method. Avoid applying this product when plants are suffering from moisture stress. If there is any doubt, apply the spray solution to a small test area of the crop or foliage to assess any undesirable effects or phytotoxicity before general application.

#### Method

[Foliar spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[Any time except during bloom.](#)