

FOLIAR APPLICATION: TREES AND VINE CROPS (EXCLUDING CITRUS)

General Information

GENERAL INFORMATION

Brandt Sequestar 14% Cu is a fully chelated, water soluble, powdered form of copper EDTA and is recommended for the prevention and correction of copper deficiency in crops. The preferred use is as a soil application to prevent the deficiency although it may also be used as a foliar spray to provide correction when a soil application is impractical.

Limitations, Restrictions, and Exceptions

Foliar Application: BRANDT SEQUESTAR 14% Cu may be applied in water or in combination with most pesticides. After the correct dilution has been made, the spray solution should be buffered to pH 6-6.5. Thorough coverage and the use of wetting agents often enhance nutrient uptake from foliar sprays. Application rates and dilution factors depend on crop sensitivity, the amount of foliage to be sprayed and the application method. Avoid applying this product when plants are suffering from moisture stress. If there is any doubt, apply the spray solution to a small test area of the crop or foliage to assess any undesirable effects or phytotoxicity before general application.

TREES & VINE CROPS

Apply 1 to 2 lbs. per acre in sufficient spray volume for thorough coverage (25 gal. minimum) at dormant/delayed dormant before bloom, then after full bloom at 2 to 3 week intervals as needed.

NOTE: For deciduous crops when fruit is present, apply 1 lb per acre in 50-100 gal of water.

Method

[Foliar spray](#)

Rates

field_rates 0

•

Timings

At dormant/delayed dormant before bloom