

TURF - TURFGRASSES FOR SOD (FL ONLY) - MUCK OR PEAT

General Information

GENERAL INFORMATION

IMPORTANT: Read the entire "DIRECTIONS FOR USE" and the "WARRANTY—CONDITIONS OF SALE" before using this product. Apply this herbicide before weeds emerge or after removal of weed growth. This product controls a wide variety of annual Broadleaf and Grass weeds when used at selective rates in agricultural crops and ornamental plantings. Where a range of application rates is given, use the low rate on Coarser textured soil and soil lower in organic matter; use the high rate on Finer textured soil and soil higher in organic matter. To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result. Since this product enters weeds mainly through their roots, moisture is needed to move it into the root zone. Very dry soil conditions and lack of rainfall following application may necessitate shallow cultivation. This product is noncorrosive to equipment, nonflammable and has low electrical conductivity. This product is a Group 5 Herbicide. Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on the label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in registered combinations or in sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered herbicide which is not solely a Group 5 Herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

Limitations, Restrictions, and Exceptions

TURF

TURFGRASSES FOR SOD (FL Only)

If weeds regrow, apply an additional 2.2 pounds on Muck or Peat, or 1.1 pounds on Sandy soil.

Use Precautions and Restrictions: (To avoid crop injury): 1) Do not apply within 30

days prior to cutting or lifting. 2) Do not apply in combination with surfactants or other spray additives. 3) Use only on Turfgrass reasonably free of infestations of insects, nematodes and diseases. 4) On newly sprigged Turfgrass, temporary slowing of growth may follow application.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[New Beds: 3 to 4 days after sprigging or plugging.](#)

[Old beds: Within 2 days after lifting of sod.](#)