

FOR USE AS A SPREADER / STICKER - AIR, GROUND OR CHEMIGATION

General Information

DIRECTIONS FOR USE

BRANDT UMBRELLA is a spreader/sticker designed to increase the efficiency of various pesticides. It forms a rainfast film which resists weathering and enhances the efficacy of pesticides. BRANDT UMBRELLA may be used on agricultural, forestry, turf and ornamental, industrial, structural and non-cropland sites. NOT FOR AQUATIC USE IN WASHINGTON.

Read and follow the precautions, restrictions and recommendations on the labels of pesticides used with BRANDT UMBRELLA. Use according to the most restrictive label directions for each product in any tank mix.

COMPATIBILITY: BRANDT UMBRELLA is compatible with most pesticides. However, if the desired combination has not been previously used, a compatibility test is recommended.

MIXING: Shake well before using. Fill spray tank $\frac{1}{2}$ full of water and begin agitation. Add the recommended amount of BRANDT UMBRELLA. Add products as directed by label or in the following sequence and continue filling tank: (1) Dry flowables or water dispersing granules, (2) Wettable powders, (3) Flowables, (4) Solutions, (5) Emulsifiable concentrates. Continue agitation until spray solution is completely mixed. Continuous agitation of finished spray is recommended. If spray solution has been allowed to stand, thoroughly agitate and remix before application.

RECOMMENDATIONS: Specific use rates will vary with conditions of application such as water hardness, application method, equipment, spray droplet size, condition of foliage, etc. Smooth, waxy, or hairy foliage will require higher rates. Also, higher rates than those below may be used if recommended by pesticide labeling. Follow pesticide label directions. However, do not add this product at a rate which exceeds 5% of the finished spray volume.

Limitations, Restrictions, and Exceptions

FOR USE AS A SPREADER/STICKER (air, ground or chemigation): Use 4-16 fl.oz. per

acre. For improved efficacy, use at least 4 fl.oz. per acre when finished spray volume is less than 20 gallons per acre.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)