

## **OTHER FRUIT**

### General Information

For use as a fertilizer in foliar and fertigation applications to supplement phosphorus, potassium, calcium and zinc levels on agricultural crops.

### Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- TruPhos MeriStim is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAg representative for application timing and necessity of its use.

Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer.

### Limitations, Restrictions, and Exceptions

The number and frequency of applications determine application rates.

Apply no more than 4 quarts/acre/application.

Apply no more than 8 gallons/season.

Always apply toward the end of an irrigation cycle.

Method

[Foliar spray](#)

[Irrigation](#)

Rates

[field\\_rates 0](#)

•

Timings

[N. A.](#)